Accepted Abbreviations

Afib, AF Atrial fibrillation

chole cholecystectomy

hys, hyst hysterectomy

PPM permanent pacemaker

PM pacemaker

M murmur (with circle)

hep hepatitis

LOC level (or loss) of consciousness, level of care

LOB loss of balance

LLD leg length discrepancy, limb length discrepancy

LL lumbar laminectomy, lid lag, left leg, left lower

schiz schizophrenia

SZ schizophrenic, seizure

Sz schizophrenia, seizure

OSA obstructive sleep apnea

OM osteomyelitis

CEA carotid endarterectomy, carcinoembryonic antigen

pneu, PN, PNM pneumonia

AN, AVN avascular necrosis

ASP, asp aspiration

asp aspirate

TR trachea

trach trachea, tracheal, tracheostomy, tracheotomy

HEM, hem hemorrhage

EF ejection fraction

V/Q ventilation/perfusion, voice quality

mast mastectomy

HD hemodialysis

MD macular degeneration, manic depression

RF rheumatic fever, respiratory failure, renal failure,

right foot

RA rheumatoid arthritis, right arm

LS, LSS lumbosacral spine

inf infarct, infection, inferior

I & D incision and drainage, irrigation and debridement,

irrigation and drainage

MCP metacarpophalangeal

UE upper extremity

LE lower extremity

DECUB, decub decubitus position, pressure ulcer

jnt , jt, j joint

BIL, bil, bi, bilat bilateral

PTH parathyroid

sp spine, speech

vol volar, voluntary, volunteer

SHLD, sh shoulder

LF left foot

LA left arm

TV tricuspid valve

MV mitral valve

AV aortic valve

exp exploratory

lap. laparotomy

chol cholesterol

TOB tobacco

PO by mouth

os mouth

MBS modified barium swallow

SP speech pathology, speech, soft palate, solid phase

S/P stand and pivot, status post

Sp speech, spine

sp. spirit, alcohol

SLP speech language pathology

HTL honey thick liquid

NTL nectar thick liquid

SS signs and symptoms

Pt, pt patient

PTA prior to admission, prior to arrival

PCA patient controlled analgesia (analgesic, anesthesia)

MED medical, medicine

N/A not applicable

NAB not at bedside

req requested, required

rep report, repeat

REV, rev review

pls please

vs. versus

inc, INC incomplete, incontinent, increase

alt alternate

TBD to be determined

Wk, wk week

mos months

BID twice daily

TID three times daily

act. activity

tol tolerance, tolerated

BR bedrest, bathroom

elev. elevated

HOB head of bed

FOB foot of bed

EOB edge of bed

res residence, resident, residue

LTD long term disability

LOS length of stay

SNF skilled nursing facility

ECF extended care facility

REC recommend

rec recommendation

WW wheeled walker

BSC bedside commode

HEP home exercise program

AMB, amb ambulate, ambulation, ambulatory

INDEP, ind, I independent

SBA stand by assist

min minimal

mod moderate

max maximum

dep dependent

W with, wife

w weight, wife

wt, w weight

func, funct function, functional

mob, mobil mobility, mobilization

asst assistant

ther ex therapeutic exercise

BAL, bal balance

GT gait, gait training, great toe

tra, trf transfer

trg, trng training

sig significant

SO significant other

WBAT weight bearing as tolerated

NWB non-weight bearing

FWB full weight bearing

TTWB touch-toe weight bearing, toe touch weight bearing

PWB partial weight bearing

R, rt, r right

L, lt, l left

st straight

H/O, h/o history of

SX, Sx signs, surgery, symptoms

D/T due to

dec decreased

deb debridement

debil debility

dehyd dehydration

MOW meals on wheels

EtOH ethyl alcohol

ED emergency department

RO, R/O rule out

PM&R physical medicine and rehabilitation

DME durable medical equipment